

# Satellite RNAs interfere with the function of viral RNA silencing suppressors

## OPEN ACCESS

### Edited by:

K. Andrew White,  
York University, Canada

### Reviewed by:

Juan Antonio Garcia,  
Consejo Superior de Investigaciones  
Científicas, Spain  
Ricardo Flores,  
Instituto de Biología Molecular y  
Celular de Plantas (UPV-CSIC), Spain

### \*Correspondence:

Chang-Yong Zhou,  
National Citrus Engineering Research  
Center, Citrus Research Institute,  
Chinese Academy of Agricultural  
Sciences, 15 Citrus Village, Xiema,  
Beibei, Chongqing 400712, China  
zhoucy@swu.edu.cn;  
Ming-Bo Wang,  
Commonwealth Scientific and  
Industrial Research Organisation Plant  
Industry, Clunies Ross Street,  
Canberra, ACT 2601, Australia  
ming-bo.wang@csiro.au

### † Present Address:

Phil Chi Khang Au,  
University of Western Australia,  
Faculty of Medicine, Dentistry and  
Health Sciences, Perth, WA, Australia

### Specialty section:

This article was submitted to  
Virology,  
a section of the journal  
Frontiers in Plant Science

**Received:** 20 January 2015

**Accepted:** 08 April 2015

**Published:** 24 April 2015

### Citation:

Shen W-X, Au PCK, Shi B-J, Smith  
NA, Dennis ES, Guo H-S, Zhou C-Y  
and Wang M-B (2015) Satellite RNAs  
interfere with the function of viral RNA  
silencing suppressors.  
Front. Plant Sci. 6:281.  
doi: 10.3389/fpls.2015.00281

Wan-Xia Shen<sup>1,2</sup>, Phil Chi Khang Au<sup>2†</sup>, Bu-Jun Shi<sup>3</sup>, Neil A. Smith<sup>2</sup>,  
Elizabeth S. Dennis<sup>2</sup>, Hui-Shan Guo<sup>4</sup>, Chang-Yong Zhou<sup>1\*</sup> and Ming-Bo Wang<sup>2\*</sup>

<sup>1</sup> National Citrus Engineering Research Center, Citrus Research Institute, Southwest University, Chongqing, China,

<sup>2</sup> Commonwealth Scientific and Industrial Research Organisation Plant Industry, Canberra, ACT, Australia, <sup>3</sup> Department of  
Plant Science, Waite Institute, Adelaide University, Glen Osmond, SA, Australia, <sup>4</sup> State Key Laboratory of Plant Genomics,  
Institute of Microbiology, Chinese Academy of Sciences, Beijing, China

Viral satellite RNAs (satRNAs) are small subviral RNAs and depend on the helper virus for replication and spread. satRNAs can attenuate helper virus-induced symptoms, the mechanism of which remains unclear. Here, we show that two virus-encoded suppressors of RNA silencing (VSRs), *Cucumber mosaic virus* (CMV) 2b and *Tombusvirus* P19, suppress hairpin RNA (hpRNA)-induced silencing of a  $\beta$ -glucuronidase (GUS) gene in *Nicotiana benthamiana*. This suppression can be overcome by CMV Y-satellite RNA (Y-Sat) via the Y-Sat-derived small interfering RNAs (siRNAs), which bind to the VSRs and displace the bound hpGUS-derived siRNAs. We also show that microRNA target gene expression in *N. tabacum* was elevated by CMV infection, presumably due to function of the 2b VSR, but this upregulation of microRNA target genes was reversed in the presence of Y-Sat. These results suggest that satRNA infection minimizes the effect of VSRs on host siRNA and microRNA-directed silencing. Our results suggest that the high abundance of satRNA-derived siRNAs contributes to symptom attenuation by binding helper virus-encoded VSRs, minimizing the capacity of the VSRs to bind host siRNA and miRNA and interfere with their function.

**Keywords:** virus-encoded suppressor, satellite RNA, symptom attenuation, RNA silencing, microRNA expression

## Introduction

SatRNAs (satRNAs) depend on their associated viruses (helper viruses) for replication, encapsidation, systemic movement and transmission, and are regarded as parasites of their helper viruses. They have small RNA genomes sized from 220 to 1400 nucleotides (nt), which share little or no sequence homology with the genome of helper viruses (Roossinck et al., 1992; Simon et al., 2004). The relatively large (800–1400 nt) satRNAs can encode one or a few proteins, but the majority of satRNAs, with <700 nt genome, possess no mRNA activity and do not encode any protein (Roossinck et al., 1992; Simon et al., 2004). Some satRNAs can induce specific symptoms in virus-infected plants (Garcia-Arenal and Palukaitis, 1999; Hu et al., 2009), for example, yellowing symptoms in *Nicotiana* species induced by Y satRNA (Y-Sat) of *Cucumber mosaic virus* (CMV) (Masuta and Takanami, 1989). Recent studies showed that the yellowing symptoms are the result of the silencing of the chlorophyll biosynthetic gene *CHLI* induced by Y-Sat-derived small interfering RNAs (siRNAs), which have a 22-nt sequence homologous to *CHLI* (Shimura et al., 2011; Smith et al., 2011). In contrast to Y-Sat, most satRNAs do not induce their own

symptoms. Another feature of satRNAs is the attenuation of helper virus-induced symptoms (Roossinck et al., 1992; Simon et al., 2004). This property has been used in transgenic plants for viral disease prevention (Baulcombe et al., 1986; Gerlach et al., 1987; Harrison et al., 1987). However, how satRNAs attenuate the symptoms remains a matter for debate. It was proposed that satRNAs might compete with the helper viruses for RNA replicase or increase antiviral silencing against the accumulation of the helper viruses (Piazzolla et al., 1982; Wu and Kaper, 1995; Hou et al., 2011). However, in some cases, satRNAs attenuate the symptoms but do not reduce the accumulation of helper viruses (Harrison et al., 1987; Moriones et al., 1992). Therefore, other unknown factors may also be involved in the regulation of helper virus-induced symptoms.

RNA silencing is a sequence-specific RNA degradation process induced by 21–24-nt small RNAs (sRNAs) processed from double-stranded RNA (dsRNA) or self-complementary hairpin RNA (hpRNA) by Dicer or Dicer-like protein, an RNase III-like endoribonuclease. One strand of each sRNA is loaded onto the RNA-induced silencing complex (RISC) and guides Argonaute (AGO) proteins in the RISC to cleave targeted RNA (Zamore, 2001). In plants RNA silencing has been recognized as an antiviral defense mechanism by which siRNAs derived from the virus guide the degradation of the viral RNA in the infected plant (Baulcombe and Molnar, 2004; Ruiz-Ferrer and Voinnet, 2009). To overcome this host defense mechanism, viruses have evolved a counter-defense strategy by encoding suppressors of RNA silencing (VSRs) (Zhang et al., 2006; Shimura and Pantaleo, 2011; Pumplin and Voinnet, 2013). A predominant action mode of VSRs is to bind the siRNAs preventing their entry into the RISC and degradation of the viral RNA (Lakatos et al., 2006). Beside viral siRNAs, VSRs also interfere with the function of host small RNAs including microRNAs (miRNAs) (Kasschau et al., 2003; Chapman et al., 2004; Zhang et al., 2006; Lewsey et al., 2007). miRNAs play a critical role in processes such as cell division, leaf formation, and flower development in plants (Voinnet, 2009). Therefore, while VSRs help the viruses to evade antiviral RNA silencing by the host, they can also induce symptoms by disrupting functions of specific miRNA species, which regulate host plant development (Zhang et al., 2006; Lewsey et al., 2007; Jay et al., 2011; Du et al., 2014).

In this study, we used an *Agrobacterium* infiltration (agro-infiltration)-based transient expression system to investigate how the satRNA of CMV Y (Y-Sat) affected the ability of CMV-encoded VSR 2b and *Tombusvirus*-encoded VSR P19 to suppress hpRNA-induced gene silencing in *Nicotiana benthamiana*. Both VSRs suppressed hpRNA-induced gene silencing, but in the presence of Y-Sat, this VSR effect was reversed. Using RNA immunoprecipitation, we showed that this reversion is due to the bound hpRNA-derived siRNAs being displaced from the VSRs by Y-Sat-derived siRNAs. We also found that Y-Sat infection minimizes the over-expression of target genes of endogenous miRNAs caused by CMV infection, indicating that Y-Sat inhibited the effect of the 2b VSR on the function of host miRNAs.

## Materials and Methods

### Plant Growth, Virus Inoculation, Plasmid Constructs, and Plant Transformation

*N. benthamiana* and *N. tabacum* (HcPro transgenic; Wang et al., 2004) plants were grown at 25°C under a 16-h light/8-h dark cycle. For virus infection, leaves of 3–5 week old plants were dusted with carborundum and rub-inoculated with leaf extracts of previously infected plants in 0.1 M phosphate buffer (pH 7.2).

The constructs of *GFP* and *P19* driven by the CaMV 35S promoter were previously described (Wood et al., 2009). The *2b* (wild-type) and *2bm* (mutant; created by substituting “C” for “U” in the start codon AUG, as well as three other AUG codons present in the *2b* coding sequence) constructs were also previously described (Hou et al., 2011). The *GUS* construct (pIG121Hm) contains an intron in the coding sequence, which was designed to prevent *GUS* expression in bacterial cells (Ohta et al., 1990; Ishikawa, 2009). The *hpGUS* construct was previously reported, designed to express a long *GUS* hairpin RNA with a 563 bp dsRNA arm and an 1113 nt loop (Wang and Waterhouse, 2000). All of the above constructs were introduced into *A. tumefaciens* strains AGL1 (*GFP* and *hpGUS*) or GV3101 (pIG121Hm, *2b*, *2bm*, and *P19*).

### *Agrobacterium* Infiltration Assay

*A. tumefaciens* strains containing plant expression constructs were grown overnight at 28°C in Luria-Bertani medium (LB) containing appropriate antibiotics. After centrifugation, *Agrobacterium* cells were re-suspended in buffer containing 10 mM MgCl<sub>2</sub> and 150 µg/mL acetosyringone, with a final optical density at 600 nm (OD<sub>600</sub>) of 0.7. The bacterial suspensions were incubated at room temperature for ~3 h and then infiltrated into expanded *N. benthamiana* leaves using flat-pointed syringe (Liu and Lomonosoff, 2002). For protein and RNA isolation, agro-infiltrated leaf sections were visualized under a blue light torch (NightSea™, DFP-1™, for exciting green light emission), excised with a pair of scissors and immediately frozen in liquid nitrogen at 4 or 5 days post agro-infiltration (dpa).

### MUG Assay of GUS Expression

For fluorometric MUG (4-methyl-umbelliferyl-β-D-galacturonide) assay, agro-infiltrated leaf sections were homogenized in extraction buffer (50 mM sodium phosphate, 10 mM Na<sub>2</sub>EDTA, 0.1% sarcosyl, 10 mM β-mercaptoethanol, pH 7.0). The homogenates were subjected to centrifugation for 5 min at 4°C at 16,000 × g, and the supernatants were collected and kept on ice. Concentration of total protein was determined using the Bradford reagent. MUG reactions were carried out at 37°C and GUS activities were calculated essentially as previously described (Chen et al., 2005) using a fluorescence plate reader (Wallac 1420 VICTOR, PerkinElmer, Turku, Finland).

## RNA Isolation, Northern Blot Analysis, and RT-PCR

Total RNA was isolated from *N. benthamiana* and *N. tabacum* leaves or leaf sections using Trizol reagent (Invitrogen) following the manufacturer's instructions. Isopropanol precipitation of the total RNA was carried out overnight at  $-20^{\circ}\text{C}$  to maximize recovery of the small RNA fraction.

For detection of large RNAs in northern blot hybridization, total RNA ( $\sim 5\text{--}10\text{ }\mu\text{g}$ ) was separated in formaldehyde-agarose gel (Sambrook et al., 1989) and blotted onto Hybond-N membrane (Amersham Life Science, NSW, Australia) in  $10\times$  SSC. The membrane was hybridized with [ $\alpha\text{-}^{32}\text{P}$ ] UTP-labeled RNA probes obtained by *in vitro* transcription using radioactive UTP from the respective sequences (Supplemental Text 1) cloned into pGEM plasmids (Promega, NSW, Australia). Hybridization was performed as previously described (Wang et al., 2001).

For detection of small RNAs in northern blot hybridization, approximately  $10\text{--}20\text{ }\mu\text{g}$  of total RNA were separated in a denaturing (urea) 17% polyacrylamide gel. RNA was electroblotted onto a Hybond-N<sup>+</sup> membrane (Amersham Life Science) in  $0.5\times$  TBE buffer, UV cross-linked, and hybridized overnight at  $42^{\circ}\text{C}$  with [ $\alpha\text{-}^{32}\text{P}$ ] UTP labeled, carbonate buffer-fragmented RNA probe (Wang et al., 2001). The hybridized membrane was washed three times (20 min per wash) at  $40^{\circ}\text{C}$  with  $2\times$  SSC containing 0.2% SDS. Both large and small RNA northern blots were visualized on a phosphorimager (Fujifilm FLA 5000, Tokyo, Japan).

For RT-PCR or qRT-PCR analysis of 2b, HcPro mRNA, or miRNA target genes, total RNA ( $\sim 1\text{ }\mu\text{g}$ ) isolated from infected and uninfected leaves was reverse-transcribed with Superscript III reverse transcriptase (Invitrogen) following the manufacturer's instruction using oligo-dT<sub>23</sub> as primer. The synthesized cDNA was amplified using 2b, HcPro or miRNA target gene-specific primers (see Supplemental Text 2 for primer sequences) with  $56^{\circ}\text{C}$  (2b) or  $60^{\circ}\text{C}$  (HcPro and miRNA target genes; qRT-PCR) as the annealing temperature. PCR cycle numbers (2b) are given on the gel pictures. Real-time PCR was performed in the Rotor-Gene 6000 (Corbett Life Science, San Francisco, USA) real-time rotary analyser using SYBR Green reagent and Platinum Taq polymerase (Invitrogen) in three technical replicates for each sample. mRNA encoding the *N. tabacum* elongation factor 1 $\alpha$  (EF1 $\alpha$ ) was amplified for use as loading control.

## RNA Immunoprecipitation

RNA immunoprecipitation was performed as described by Omarov et al. (2006) with minor changes. Briefly, tissues (2g) were ground into fine powder in liquid nitrogen and re-suspended in 2 mL ice-cold extraction buffer with 150 mM HEPES (pH 7.5), 200 mM NaCl, 1 mM EDTA, 2 mM DTT, 16 unit RNaseOUT (Invitrogen, Victoria, Australia) and 20  $\mu\text{L}$  Plant protease inhibitors (Sigma-Aldrich, NSW, Australia). The homogenized plant material was filtered through two layers of miracloth (Calbiochem, Victoria, Australia) and centrifuged twice at  $10,000\times g$  at  $4^{\circ}\text{C}$  for 15 min. The supernatant was pre-cleared in 10  $\mu\text{L}$  Salmon Sperm DNA/Protein G agarose beads (Millipore, Victoria, Australia) for 30 min at  $4^{\circ}\text{C}$  under constant rotation. An aliquot (800  $\mu\text{L}$ ) of pre-cleared lysate was

then immunoprecipitated with 2  $\mu\text{L}$  of P19 rabbit antibodies (A gift from Prof. Herman Bertus Scholthof, Texas A&M University) and 30  $\mu\text{L}$  of Salmon Sperm DNA/Protein G agarose beads (Millipore) for 3 h at  $4^{\circ}\text{C}$  under constant rotation. Following immunoprecipitation, the agarose-antibody-P19 /RNA complexes were washed six times in extraction buffer. P19/RNA complexes were eluted with 100  $\mu\text{L}$  of RNA-IP elution buffer [100 mM Tris-HCl pH 8.0, 10 mM EDTA, 1% SDS, 40 units of RNaseOUT (Invitrogen), 20  $\mu\text{g}$  of Proteinase K (Sigma-Aldrich)] at  $65^{\circ}\text{C}$  for 1 h. RNA was purified by extraction with an equal volume of acidic phenol/chloroform (Ambion, Applied Biosystems, Victoria, Australia), followed by overnight ethanol precipitation in the presence of acidic sodium acetate and 20  $\mu\text{g}$  glycogen (Fermentas, Thermo fisher scientific, Victoria, Australia) at  $-80^{\circ}\text{C}$ , and resuspended in DEPC-treated H<sub>2</sub>O for small RNA northern blot hybridization. For Western blot analysis of the pull-down material, elution step was omitted and washed beads were stored at  $-20^{\circ}\text{C}$  for Western blotting.

## Western Blot Analysis

Beads containing the immunoprecipitated P19/RNA complexes were denatured at  $95^{\circ}\text{C}$  for 5 min with addition of Western blot loading buffer (Bromophenol blue, 0.125 M Tris-HCl, 4% SDS, 20% glycerol, 2%  $\beta$ -mercaptoethanol, pH 6.8). The proteins were separated on a 15% SDS polyacrylamide gel, transferred to Immobilon P membrane (Millipore) and detected by chemiluminescence using 1/2500 anti-P19 antibody (Omarov et al., 2006) and 1/5000 anti-rabbit Ig HRP conjugate (Chemicon, Victoria, Australia).

## Results


### Y-Sat Reduces the Symptoms Caused by the Helper Virus Fny-CMV

The strains of *Cucumber mosaic virus* (CMV) are classified into two subgroups, I and II, based on their serology, symptomatology and sequence homology (Palukaitis et al., 1992). Fny-CMV is a subgroup I strain and induces severe symptoms in many host plant species (Zhang et al., 1994). Its encoded VSR 2b has been shown to be a strong virulence contributor (Shi et al., 2002). In *N. benthamiana*, Fny-CMV not only induced severe symptoms, but also constrained plant growth (Figure 1A). However, in the presence of Y-Sat, the severity of the Fny-CMV-caused symptoms was reduced and plant growth improved, despite the bright yellow symptoms typical of Y-Sat infection (Figure 1A). This demonstrates that Y-Sat attenuates the symptoms induced by Fny-CMV, which is consistent with previous reports (Hou et al., 2011). Northern blot hybridization showed that Y-Sat co-infection reduced the level of CMV RNA, but not dramatically (Figure 1B).

### Y-Sat is the Dominant Source of Viral siRNAs in Infected Plants

Previous studies by northern blotting and small-scale sequencing showed that the replication of Y-Sat is accompanied by high levels of sRNAs derived from Y-Sat (Wang et al., 2004; Ebhardt and Unrau, 2009). To further demonstrate this, we obtained





two sets of sRNA sequence data from *N. tabacum* plants co-infected with Q-CMV (a subgroup II CMV) and Y-Sat using Illumina deep sequencing technology, and analyzed the relative abundance of Y-Sat and CMV-derived sRNAs. A total of 195,291 and 330,965 reads of 20–24 nt sRNAs (from a total of approximately 8 million sRNA reads each) were perfectly mapped to the Y-Sat genome in the two samples S1 and S2, respectively (Figure 1C). These numbers are approximately 2 and 3 times as much as the total 20–24 nt sRNA reads perfectly mapped to the Q-CMV genome (95,073 and 103,443 for S1 and S2, respectively) (Figure 1C). This is despite the 23 fold bigger size of the CMV genome (8617 nt) than the Y-Sat genome (369 nt), indicating that the Y-Sat genome has approximately 50 fold more sRNA coverage than the CMV genome, which is consistent with the previously reported small-scale sequencing result (Ebhardt and Unrau, 2009). This relatively high abundance of Y-Sat sRNAs could result from a relatively high proportion of Y-Sat RNA in the total viral RNA population of the Q-CMV+Y-Sat-infected *N. tabacum* plants, but a previous report showed that Y-Sat RNA represents about 31–38% of the total viral RNAs (Takanami, 1981), well below the 2–3 fold level of Y-Sat sRNAs.

### Y-Sat RNA Interferes with the Function of 2b in *Nicotiana benthamiana* Leaves

Cucumoviral 2b proteins are essential for cucumoviruses to cause viral symptoms (Ding et al., 1996; Shi et al., 2002; Soards et al., 2002; Zhang et al., 2006; Lewsey et al., 2007), while Y-Sat infection reduces the symptoms (Figure 1). This led us to investigate if Y-Sat infection interferes with the 2b functions. We developed an Agrobacterium infiltration (agro-infiltration)-based transient expression system. In this system equal amounts of VSR-expressing constructs were agro-infiltrated into *N. benthamiana* leaves with infection of Q-CMV or Q-CMV plus Y-Sat, and mock-treated plant leaves, and the level of hpRNA-induced silencing of the  $\beta$ -glucuronidase (GUS) reporter gene in the leaves was measured. We used the subgroup II CMV strain, Q-CMV, as the helper virus to infect *N. benthamiana* instead of Fny-CMV. This was because Q-CMV encodes a relatively weak 2b VSR (Chapman et al., 2004; Zhang et al., 2006; Ye et al., 2009) and the level of VSR activity from the infecting Q-CMV is negligible compared to that from the agro-infiltrated VSR constructs expressing a strong VSR (SD-CMV 2b and tombusvirus P19) at relatively high levels. This allowed us to examine the effect of Y-Sat specifically on the agro-infiltrated VSR. In addition, Q-CMV only induces mild symptoms in *N. benthamiana* and allowed the normal leaf expansion required for agro-infiltration. Furthermore, Q-CMV, like Fny-CMV, supports high levels of Y-Sat replication and Y-Sat siRNA accumulation (Figure 1C and Wang et al., 2004). The following constructs were used in the agro-infiltration assay: GUS reporter gene (*GUS*), GUS silencing-inducing hpRNA (*hpGUS*), strong 2b (*2b*) VSR encoded by the subgroup I Shan-Dong (SD)-CMV (Hou et al., 2011), and green fluorescent protein gene (*GFP*) as a visual marker. A construct expressing a mutant 2b gene (*2bm*), in which the start codon and three other in-frame AUG codons of the 2b coding sequence were mutated (Duan et al., 2012), was used as a negative control. Gene expression in all of these constructs was

driven by a constitutive *Cauliflower mosaic virus* (CaMV) 35S promoter.


As shown by the fluorometric MUG (4-methylumbelliferyl- $\beta$ -D-glucuronide) assay in Figure 2A, without *2b* co-infiltration, strong GUS silencing was induced by *hpGUS* in all uninfected (mock), Q-CMV-infected (Q-CMV), and Q-CMV+Y-Sat co-infected plants. Northern blotting confirmed that the *GUS* gene was silenced at the mRNA level (Figure 2B, top two panels). When *GUS* and *hpGUS* were co-infiltrated with *2bm*, *GUS* silencing was not suppressed (Figures 2A,B). However, when *GUS* and *hpGUS* were co-infiltrated with *2b*, the *GUS* activity was recovered substantially in the mock and Q-CMV-infected plants, but this was not the case for Q-CMV+Y-Sat co-infected plants, which continued to show strong GUS silencing (Figures 2A,B). These results indicate that *2b* possesses the ability to suppress the hpRNA-induced GUS silencing but this ability is greatly inhibited in the presence of Y-Sat. There were variations in CMV RNA3 levels among the different samples, with a few samples containing particularly high amounts of CMV RNA3, although the four *GUS*+*hpGUS*+*2b*-infiltrated samples showed relatively uniform levels (Figure 2B). The variations could result from different age or infection state of the leaf tissues harvested because they did not seem to be due to the presence or absence of Y-Sat. CMV-infected samples did not show enhanced GUS expression in comparison to mock-treated ones (Figure 2A), and the variations in CMV RNA3 level did not have a clear effect on GUS expression levels (Figure 2B), presumably because of the weak VSR activity of the Q-CMV 2b.

Northern blot analysis of sRNAs corresponding to the 563-bp dsRNA arm region of the *hpGUS* transcript (Wang and Waterhouse, 2000) showed that Y-Sat did not increase the accumulation of *hpGUS*-derived siRNAs in the absence (Figure 2C, lanes 3–6) or presence (lanes 7–10) of *2b*. This indicates that the enhanced GUS silencing upon Y-Sat infection was not caused by an increase in the amount of *hpGUS*-derived siRNAs. In addition, a semi-quantitative RT-PCR analysis showed that the expression level of the agro-infiltrated *2b* gene was not affected by Q-CMV or Q-CMV+Y-Sat infection (Supplemental Figure S1). These results indicate that Y-Sat minimized the suppression of the hpRNA-induced GUS silencing mediated by *2b*.

Surprisingly, when a probe complementary to the loop of the hpGUS RNA was used, an intense band of the predicted loop size (~1.1 kb) was detected in the *2b*-infiltrated tissues (Figure 2B, 2nd panel; lanes 13–16), indicating that the *2b* protein enhanced the accumulation of the hpRNA loop. Loop fragments have been known to accumulate in hpRNA-expressing plants (Wang et al., 2008). However, the intensity of the hybridized loop fragment from the tissues infected with Q-CMV+Y-Sat was much weaker than that from the tissues infected with Q-CMV alone (lanes 15–16 vs. lanes 13–14). This further suggests that Y-Sat interferes with *2b* function.

### Y-Sat RNA Interferes with VSR P19 Function in *N. benthamiana* Leaves

The tombusvirus-encoded P19 is a well-characterized VSR that suppresses RNA silencing via binding and sequestering sRNAs (Vargason et al., 2003; Baulcombe and Molnar, 2004; Lakatos



**FIGURE 2 | The suppression of hpRNA-induced GUS silencing by agro-infiltrated 2b is released by CMV+Y-Sat infection. (A)** MUG assay of GUS expression in Mock (inoculated only with buffer), Q-CMV and Q-CMV+Y-Sat-infected *N. benthamiana* leaves infiltrated with the constructs indicated. The plants were infiltrated at 11 or 13 days after inoculation. The GUS activity was analyzed at 4 days post agroinfiltration (dpa). **(B)** Northern blot hybridization analysis of GUS expression in a subset of the samples described in (A). Ten microgram of total RNA from each sample was

hybridized with an antisense GUS probe to detect GUS mRNA (top panel), a probe complementary to the loop of the hpGUS RNA to detect both GUS mRNA and hpGUS loop fragments (second panel), a CMV RNA 3a probe to detect CMV (third panel), and a Y-Sat probe to detect Y-Sat (fourth panel) in the agroinfiltrated tissues. Ethidium bromide-stained rRNA bands are used as a loading control (bottom panel). **(C)** Northern blot hybridization analysis of small RNAs in a subset of the samples described in (A). Twenty microgram (Continued)



**FIGURE 2 | Continued**

of total RNA each was hybridized with a RNA probe corresponding to the dsRNA stem region of *hpGUS* to detect *hpGUS*-derived siRNA (top panel), an antisense Y-Sat RNA probe to detect Y-Sat-derived siRNAs (middle panel) and a U6 probe to show loading control (bottom panel).

The 21 and 24-nt siRNA size markers are loaded in the last lane on the right. As can be seen, the level of *hpGUS*-derived siRNAs is not significantly affected by Q-CMV Y-Sat infection. SD2b or SD2bm, wild-type or mutant 2b derived from the group I Shandong (SD) strain of CMV.

et al., 2006). We examined if Y-Sat also interferes with P19 functions. The same agro-infiltration system used above for 2b was applied. P19 suppressed *hpGUS*-induced GUS silencing in mock and Q-CMV-infected *N. benthamiana* tissues, and the suppression was negated in Q-CMV+Y-Sat-infected tissues (**Figures 3A,B**). In addition, the presence of Y-Sat did not increase the accumulation of *hpGUS*-derived siRNAs (**Figure 3C**), indicating that the recovery of GUS silencing in Q-CMV+Y-Sat-infected tissues was not due to an increase in the amount of *hpGUS*-derived siRNAs. These results indicate that Y-Sat enhances *hpRNA*-induced silencing in the agro-infiltrated *N. benthamiana* leaves by interfering with the function of P19 minimizing P19-mediated suppression of silencing. Thus, Y-Sat may use the same mechanism to interfere with the functions of 2b and P19.

It is worth mentioning that both 2b and P19 enhanced GUS expression in the *GUS*-infiltrated plants lacking *hpGUS* (**Figures 2A,B** and **Figures 3A,B**; compare “GUS” or “GUS+2bm” with “GUS+2b” or “GUS+P19”). This probably resulted from VSR-mediated suppression of sense co-suppression or posttranscriptional gene silencing that occurs to agro-infiltrated transgenes in *N. benthamiana* leaves (Voinnet et al., 1999). In the presence of Y-Sat, part of the enhancement of GUS expression by 2b or P19 was lost (**Figures 2B, 3B**; compare lanes 11–12 with lanes 9–10), further supporting the conclusion that Y-Sat interferes with the function of 2b and P19.

### Y-Sat Infection Minimizes the Effect of CMV Infection on Expression of miRNA Target Genes


VSRs, including CMV 2b, suppress the function of both viral and host siRNAs and miRNAs (Brigneti et al., 1998; Kasschau et al., 2003; Pumplin and Voinnet, 2013). Our agro-infiltration assays indicated that Y-Sat infection interferes with the VSR effect on siRNA-directed gene silencing in the host plant. To determine if Y-Sat infection affected the function of VSRs on host miRNA-directed gene silencing, expression levels of target genes of five miRNAs (miR156, miR160, miR168, miR172, and miR395) in the Q-CMV and Q-CMV+Y-Sat-infected transgenic *N. tabacum* plants overexpressing potyvirus (*Tobacco etch virus*)-encoded VSR HcPro were analyzed. The HcPro plants rather than wild-type plants were used for two reasons. First, the plants do not develop yellowing symptoms with Q-CMV+Y-Sat infection, presumably because Y-Sat siRNA-directed silencing of the chlorophyll biosynthetic gene *CHL1* is repressed by HcPro (Shimura et al., 2011; Smith et al., 2011; Shen et al., 2014). This minimizes the possible effect of the yellowing symptoms on the expression of miRNAs' targets. Second, infection of the HcPro plants, but not wild-type plants, with Q-CMV resulted in consistent upregulation of miRNA target gene expression (**Figure 4A** and data not shown), possibly because of a synergistic effect between CMV 2b

and HcPro or the relatively high level of CMV accumulation in HcPro plants in comparison to wild-type plants. This allowed us to examine the effect of Y-Sat on VSR-mediated mis-regulation of miRNA target genes. It is worth noting that HcPro increases the expression levels of several miRNAs' targets in *Arabidopsis* (Kasschau et al., 2003), but among the five *N. tabacum* miRNA target genes examined, only that of miR160 appeared to be upregulated in the HcPro-expressing *N. tabacum* plants without CMV infection (**Figure 4A**). The target genes of the remaining four miRNAs were expressed at similar levels between HcPro-expressing (HcPro+) (samples 3–4) and wild-type (HcPro-) (samples 1–2) *N. tabacum* plants (**Figure 4A**).

All five miRNA target genes showed increased expression in the CMV-infected HcPro+ plants compared to the uninfected HcPro+ plants (**Figure 4A**, samples 5–6 vs. 3–4). This suggests that the 2b protein from the infecting CMV, probably in conjunction with the transgene-expressed HcPro, suppressed miRNA-directed silencing resulting in upregulation of the miRNA target genes in the CMV-infected plants. However, in the plants co-infected with CMV and Y-Sat, the expression levels of these target genes were reduced and became similar to those in the uninfected HcPro+ plants (compare samples 7–8 with samples 5–6 or 3–4). This change was not due to a change in HcPro expression because its expression level was similar between the CMV and CMV+Y-Sat-infected plants and showed no correlation with the expression of miRNA target genes (**Figure 4B**). Furthermore, this change was not caused by a change in CMV RNA level because there was in general no correlation between the expression level of miRNA target genes and the amount of CMV RNA in the CMV and CMV+Y-Sat-infected plants (**Figures 4A,C**). The expression of the miR156 target gene in the two CMV-infected plants appeared to be an exception, which showed a positive correlation with the amount of CMV RNA (**Figures 4A,C**). Taken together, the decrease in miRNA target gene expression upon Y-Sat infection suggests that the suppression of miRNA function by CMV 2b in conjunction with HcPro was overcome by Y-Sat.

### Y-Sat siRNAs Bind to P19 and Sequester P19 from *hpGUS*-Derived siRNAs


As shown in **Figure 1**, Y-Sat-derived siRNAs were highly abundant in the infected plants. This led us to test if the satRNA-derived siRNAs (sat-siRNAs) might bind and sequester the VSRs. RNA immunoprecipitation (RNA-IP) was performed on the *N. benthamiana* tissues agro-infiltrated with *GUS*+*hpGUS*+P19 using the P19 antibody ( $\alpha$ -P19). Northern analysis of the P19-bound siRNAs showed that Y-Sat-siRNAs were present at high levels in the  $\alpha$ -P19 RNA-IP from the Q-CMV+Y-Sat-infected tissues (**Figure 5A**, middle-right panel). In contrast, the amount of *hpGUS*-derived siRNAs in the  $\alpha$ -P19 RNA-IP was dramatically reduced compared to that in the plants



**FIGURE 3 | The suppression of hpRNA-induced GUS silencing by agro-infiltrated P19 is released by CMV+Y-Sat infection. (A)** MUG assay of GUS expression in Mock, Q-CMV and Q-CMV+Y-Sat-infected (17 days post inoculation) *N. benthamiana* leaves infiltrated with the indicated constructs at 5 dpa. **(B)** Northern blot hybridization analysis of GUS expression in a subset of the samples described in **(A)**. Five microgram of total RNA was hybridized with an antisense GUS probe to detect GUS mRNA levels (top panel), a CMV RNA 3a probe to measure CMV levels (second panel) and a Y-Sat probe to measure Y-Sat levels

(third panel) in the agroinfiltrated tissues. Ethidium bromide-stained rRNA bands were used as loading control (bottom panel). **(C)** Northern blot hybridization analysis of small RNAs in a subset of the samples described in **(A)**. Ten microgram of total RNA each was hybridized with a RNA probe corresponding to the dsRNA stem region of hpGUS to detect hpGUS-derived siRNA (top panel) and a U6 probe to show loading control (bottom panel). The sizes of the detected siRNAs are indicated. As can be seen, the level of hpGUS-derived siRNAs is not significantly affected by Q-CMV Y-Sat infection.






**FIGURE 4 | The increased expression of miRNA target genes by CMV infection in HcPro-expressing *N. tabacum* plants is reversed in the presence Y-Sat. (A)** qRT-PCR analysis of predicted target genes for miR156, miR160, miR168, miR172, and miR395. The EF1 $\alpha$  gene was used as the internal reference. **(B)** qRT-PCR analysis of

HcPro expression in the HcPro plants. **(C)** Northern blot analysis of Q-CMV and Y-Sat levels in the transgenic HcPro *N. tabacum* plants used for qRT-PCR analysis. HcPro+, *N. tabacum* plants containing the HcPro transgene; HcPro-, *N. tabacum* segregants containing no HcPro transgene.

infected by Q-CMV alone without Y-Sat (Figure 5A, top-right panel and Figure 5B). These results indicate that P19 suppresses the silencing of *GUS* expression via binding to the *hpGUS*-derived siRNAs, and that in the presence of Y-Sat, the abundant Y-Sat-derived siRNAs bind to P19 reduce its capacity to bind *hpGUS*-derived siRNAs. As a result, Y-Sat

infection allowed more *hpGUS*-derived siRNAs to be available for directing *GUS* silencing. We did not pursue an RNA-IP assay for the binding of Y-Sat siRNAs to VSR 2b due to the unavailability of 2b antibody. However, from the previous studies showing that 2b also suppresses RNA silencing by binding siRNAs (Vargason et al., 2003; Baulcombe and Molnar,



2004; Gonzalez et al., 2012), it is expected that the abundant Y-Sat-derived siRNAs would bind to 2b and reduces the amount of 2b-bound *hpGUS*-derived siRNAs, enhancing GUS silencing in the *N. benthamiana* tissues agro-infiltrated with *GUS*+*hpGUS*+2b.

## Discussion

A previous study on defective interfering RNAs (DI-RNAs) associated with Tomato bushy stunt tobravirus (TBSV) has shown that abundant siRNAs derived from the DI-RNAs can saturate TBSV-encoded P19 VSR and elevate the levels of free TBSV-derived siRNAs (Havelda et al., 2005). These free TBSV-derived siRNAs in the presence of the DI-RNAs have been proposed to play a role in directing silencing against the helper virus, thereby reducing TBSV accumulation and symptom severity (Havelda et al., 2005). A recent study on a CMV satRNA has also implicated increased anti-helper virus RNA silencing and reduced helper

virus accumulation in satRNA-mediated symptom attenuation (Hou et al., 2011). This study shows that satRNA reduces the accumulation of VSR 2b. Because 2b interferes with the function of viral sRNAs, it is likely that satRNA enhances anti-viral silencing (Hou et al., 2011) which in turn reduces the accumulation of CMV helper virus.

Whereas increased antiviral silencing and reduced helper virus accumulation as proposed in the two studies may well play a role in satRNA-mediated symptom attenuation, our results suggest an alternative mechanism by which satRNAs reduce helper virus-caused symptoms. Similar to the DI RNA scenario, satRNA-derived siRNAs bind to and saturate VSRs due to their high abundance in virus-infected tissues. However, in addition to promoting RNA silencing against the helper virus to reduce its accumulation, this minimizes the capacity of VSRs to bind host endogenous siRNAs and miRNAs. As a result, these endogenous siRNAs and miRNAs are free to form RISC with AGO proteins allowing normal sRNA-mediated gene regulation to occur in the host plants. Thus, developmental abnormalities, or symptoms, caused by the helper virus VSRs are minimized in the presence of satRNAs.

We have demonstrated that Y-Sat effectively reduced the ability of 2b and P19 in suppressing the *hpRNA*-induced GUS silencing in *N. benthamiana* plants using the agro-infiltration-based transient expression assay. We have also shown that CMV infection increased the expression levels of miRNAs' target genes but in the presence of Y-Sat the expression levels of these target genes reverted to uninfected levels. Taken together, our data indicate that Y-Sat can impact on both the siRNA- and miRNA-directed RNA silencing pathways in the host plants. CMV 2b was previously shown to bind both siRNAs and AGO proteins (Zhang et al., 2006; Duan et al., 2012; Hamera et al., 2012). However, *in vivo* suppression of RNA silencing requires its siRNA binding activity but is independent of its ability to interact directly with AGOs (Duan et al., 2012). We thus expect that binding and saturation of the 2b VSR by sat-siRNAs minimizes the ability of 2b to bind host sRNAs and suppress host sRNA-directed RNA silencing, which would restore host sRNA-mediated gene regulation that would benefit normal plant development resulting in attenuated viral symptoms.

It is interesting to note that CMV infection alone seemed to affect the function of the agro-infiltrated 2b and P19 VSRs. There is a reduction in GUS expression in the CMV-infected in comparison to uninfected plants in *GUS*+*hpGUS*+2b-infiltrated (Figure 2A) as well as *GUS*+*hpGUS*+P19-infiltrated (Figure 3A) tissues. A possible explanation is that CMV-derived siRNAs, although less abundant than Y-Sat-derived siRNAs, also bind to the VSRs reducing their capacity to suppress *hpRNA*-directed GUS silencing. This result could imply that viral siRNAs not only play a role in directing silencing against the viruses, but also contribute to the minimization of VSRs' interference with host plant sRNA functions.

satRNA-derived siRNAs have the potential to compete with host plant-encoded sRNAs for endogenous RNA silencing machineries such as DCLs and AGO proteins, which in turn could affect host sRNA-directed gene regulation. Thus, the high

abundance of satRNA siRNAs could both reduce symptoms by binding to VSRs and exacerbate symptoms by competing for host silencing machineries. However, we have shown here and previously (Shen et al., 2014) that the high abundance of Y-Sat siRNAs does not seem to affect the biogenesis of host sRNAs in *Nicotiana* plants. Furthermore, in the absence of a strong VSR and visible helper virus-caused symptoms (namely in the case of the infection of *N. benthamiana* with Q-CMV which encodes a weak 2b), host sRNA-directed silencing and plant growth are not affected by the high level of Y-Sat siRNAs (Shen et al., 2014). Why host silencing machineries are not saturated by the abundant satRNA siRNAs remains to be further understood, but these results suggest that satRNA-derived siRNAs affect host sRNA functions mainly through interacting with VSRs.

In conclusion, our results suggest that satRNAs are likely to attenuate the symptoms via their abundant siRNAs, which bind helper virus-encoded VSRs minimizing their interference with the endogenous siRNA and miRNA pathways as well as the antiviral silencing pathways as proposed in the previous studies. This model could apply to other viral or subviral RNAs that give rise to high levels of siRNAs in infected host cells, for example, DI RNAs. DI RNAs are a truncated form of their associated helper viral genomes and, like satRNAs, also depend on the helper virus for replication and can also attenuate helper virus-induced symptoms (Simon et al., 2004). Unlike satRNAs that occur mostly in plants, DI RNAs are widespread and associated with both plant viruses and animal viruses. Because animal viruses also encode VSRs such as the B2 protein encoded by Flock House virus (FHV) (Aliyari et al., 2008), it would be valuable to examine if DI RNAs modulate animal viral symptoms through a similar mechanism.

## References

- Aliyari, R., Wu, Q. F., Li, H. W., Wang, X. H., Li, F., Green, L. D., et al. (2008). Mechanism of induction and suppression of antiviral immunity directed by virus-derived small RNAs in *Drosophila*. *Cell Host Microbe* 4, 387–397. doi: 10.1016/j.chom.2008.09.001
- Baulcombe, D. C., and Molnar, A. (2004). Crystal structure of p19-a universal suppressor of RNA silencing. *Trends Biochem. Sci.* 29, 279–281. doi: 10.1016/j.tibs.2004.04.007
- Baulcombe, D. C., Saunders, G. R., Bevan, M. W., Mayo, M. A., and Harrison, B. D. (1986). Expression of biologically-active viral satellite RNA from the nuclear genome of transformed plants. *Nature* 321, 446–449. doi: 10.1038/321446a0
- Brigneti, G., Voinnet, O., Li, W. X., Ji, L. H., Ding, S. W., and Baulcombe, D. C. (1998). Viral pathogenicity determinants are suppressors of transgene silencing in *Nicotiana benthamiana*. *EMBO J.* 17, 6739–6746. doi: 10.1093/emboj/17.22.6739
- Chapman, E. J., Prokhnevsky, A. I., Gopinath, K., Dolja, V. V., and Carrington, J. C. (2004). Viral RNA silencing suppressors inhibit the microRNA pathway at an intermediate step. *Gene Dev.* 18, 1179–1186. doi: 10.1101/gad.1201204
- Chen, S., Helliwell, C. A., Wu, L. M., Dennis, E. S., Upadhyaya, N. M., Zhang, R., et al. (2005). A novel T-DNA vector design for selection of transgenic lines with simple transgene integration and stable transgene expression. *Funct. Plant Biol.* 32, 671–681. doi: 10.1071/FP05072
- Ding, S. W., Shi, B. J., Li, W. X., and Symons, R. H. (1996). An interspecies hybrid RNA virus is significantly more virulent than either parental virus. *Proc. Natl. Acad. Sci. U.S.A.* 93, 7470–7474. doi: 10.1073/pnas.93.15.7470
- Du, Z. Y., Chen, A. Z., Chen, W. H., Liao, Q. S., Zhang, H. M., Bao, Y. M., et al. (2014). Nuclear-cytoplasmic partitioning of *Cucumber mosaic virus* protein 2b determines the balance between its roles as a virulence determinant and an RNA-silencing suppressor. *J. Virol.* 88, 5228–5241. doi: 10.1128/JVI.00284-14
- If so, potential DI RNA-related strategies could be developed against animal viral diseases in the future.
- ## Funding
- WS was funded by a China Scholarship Council scholarship, grants from MOA's Public Benefit Research Foundation of China (201203076) and Chongqing Postdoctoral Science Foundation (Rc201423) and the Program for Changjiang Scholars and Innovative Research Team in University (IRT0976), and MW was supported by an Australian Research Council Future Fellowship (FT0991956).
- ## Acknowledgments
- We would like to thank Craig Wood and Christopher Cazonelli for helpful discussion and technical advice on agroinfiltration assays, Herman Bertus Scholthof for P19 antibody, Chikara Masuta for the Y-Sat infectious clone, Kenzo Nakamura for the GUS construct pIG121Hm, Vickie Vance, and Peter Waterhouse for the HcPro transgenic plants, Carl Davies for photography, and Zhiyou Du for critical reading of the manuscript. We also thank Joanne Lee for mapping the CMV and Y-Sat sRNAs.
- ## Supplementary Material
- The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fpls.2015.00281/abstract>
- Duan, C. G., Fang, Y. Y., Zhou, B. J., Zhao, J. H., Hou, W. N., Zhu, H., et al. (2012). Suppression of Arabidopsis ARGONAUTE1-mediated slicing, transgene-induced RNA silencing, and DNA methylation by distinct domains of the *Cucumber mosaic virus* 2b protein. *Plant Cell* 24, 259–274. doi: 10.1105/tpc.111.092718
- Ebbhardt, H. A., and Unrau, P. J. (2009). Characterizing multiple exogenous and endogenous small RNA populations in parallel with subfemtomolar sensitivity using a streptavidin gel-shift assay. *RNA* 15, 724–731. doi: 10.1261/rna.1235109
- Garcia-Arenal, F., and Palukaitis, P. (1999). Structure and functional relationships of satellite RNAs of *Cucumber mosaic virus*. *Curr. Top. Microbiol.* 239, 37–63. doi: 10.1007/978-3-662-09796-0\_3
- Gerlach, W. L., Llewellyn, D., and Haseloff, J. (1987). Construction of a plant disease resistance gene from the satellite RNA of *Tobacco ringspot virus*. *Nature* 328, 802–805. doi: 10.1038/328802a0
- Gonzalez, I., Rakitina, D., Semashko, M., Taliansky, M., Praveen, S., Palukaitis, P., et al. (2012). RNA binding is more critical to the suppression of silencing function of *Cucumber mosaic virus* 2b protein than nuclear localization. *RNA* 18, 771–782. doi: 10.1261/rna.031260.111
- Hamera, S., Song, X. G., Su, L., Chen, X. Y., and Fang, R. X. (2012). *Cucumber mosaic virus* suppressor 2b binds to AGO4-related small RNAs and impairs AGO4 activities. *Plant J.* 69, 104–115. doi: 10.1111/j.1365-3113X.2011.04774.x
- Harrison, B. D., Mayo, M. A., and Baulcombe, D. C. (1987). Virus-resistance in transgenic plants that express *Cucumber mosaic virus* satellite RNA. *Nature* 328, 799–802. doi: 10.1038/328799a0
- Havelda, Z., Hornyik, C., Valoczi, A., and Burgyan, J. (2005). Defective interfering RNA hinders the activity of a Tombusvirus-encoded posttranscriptional gene silencing suppressor. *J. Virol.* 79, 450–457. doi: 10.1128/JVI.79.1.450-457.2005
- Hou, W. N., Duan, C. G., Fang, R. X., Zhou, X. Y., and Guo, H. S. (2011). Satellite RNA reduces expression of the 2b suppressor protein resulting in the



- attenuation of symptoms caused by *Cucumber mosaic virus* infection. *Mol. Plant Pathol.* 12, 595–605. doi: 10.1111/j.1364-3703.2010.00696.x
- Hu, C. C., Hsu, Y. H., and Lin, N. S. (2009). Satellite RNAs and satellite viruses of plants. *Viruses* 1, 1325–1350. doi: 10.3390/v1031325
- Ishikawa, A. (2009). The Arabidopsis G-Protein beta-subunit is required for defense response against *Agrobacterium tumefaciens*. *Biosci. Biotechnol. Biochem.* 73, 47–52. doi: 10.1271/bbb.80449
- Jay, F., Wang, Y., Yu, A., Taconnat, L., Pelletier, S., Colot, V., et al. (2011). Misregulation of AUXIN RESPONSE FACTOR 8 underlies the developmental abnormalities caused by three distinct viral silencing suppressors in Arabidopsis. *PLoS Pathog.* 7: e1002035. doi: 10.1371/journal.ppat.1002035
- Kasschau, K. D., Xie, Z., Allen, E., Llave, C., Chapman, E. J., Krizan, K. A., et al. (2003). P1/HC-Pro, a viral suppressor of RNA silencing, interferes with Arabidopsis development and miRNA function. *Dev. Cell* 4, 205–217. doi: 10.1016/S1534-5807(03)00025-X
- Lakatos, L., Csorba, T., Pantaleo, V., Chapman, E. J., Carrington, J. C., Liu, Y. P., et al. (2006). Small RNA binding is a common strategy to suppress RNA silencing by several viral suppressors. *EMBO J.* 25, 2768–2780. doi: 10.1038/sj.emboj.7601164
- Lewsey, M., Robertson, F. C., Canto, T., Palukaitis, P., and Carr, J. P. (2007). Selective targeting of miRNA-regulated plant development by a viral counter-silencing protein. *Plant J.* 50, 240–252. doi: 10.1111/j.1365-3113.2007.03042.x
- Liu, L., and Lomonosoff, G. P. (2002). Agroinfection as a rapid method for propagating *Cowpea mosaic virus*-based constructs. *J. Virol. Methods* 105, 343–348. doi: 10.1016/S0166-0934(02)00121-0
- Masuta, C., and Takanami, Y. (1989). Determination of sequence and structural requirements for pathogenicity of a *Cucumber mosaic virus* satellite RNA (Y-satRNA). *Plant Cell* 1, 1165–1173. doi: 10.1105/tpc.1.12.1165
- Moriones, E., Diaz, I., Rodriguezcerez, E., Fraile, A., and Garciaarenal, F. (1992). Differential interactions among strains of *Tomato aspermy virus* and satellite RNAs of *Cucumber mosaic virus*. *Virology* 186, 475–480. doi: 10.1016/0042-6822(92)90012-E
- Ohta, S., Mita, S., Hattori, T., and Nakamura, K. (1990). Construction and expression in Tobacco of a  $\beta$ -Glucuronidase (GUS) reporter gene containing an intron within the coding sequence. *Plant Cell Physiol.* 31, 805–813.
- Omarov, R., Sparks, K., Smith, L., Zindovic, J., and Scholthof, H. B. (2006). Biological relevance of a stable biochemical interaction between the Tombusvirus-encoded P19 and short interfering RNAs. *J. Virol.* 80, 3000–3008. doi: 10.1128/JVI.80.6.3000-3008.2006
- Palukaitis, P., Roossinck, M. J., Dietzgen, R. G., and Francki, R. I. B. (1992). *Cucumber mosaic virus*. *Adv. Virus Res.* 41, 281–348. doi: 10.1016/S0065-3527(08)60039-1
- Piazzolla, P., Tousignant, M. E., and Kaper, J. M. (1982). *Cucumber mosaic virus*-associated RNA 5 IX. The overtaking of viral RNA synthesis by CARNA 5 and dsCARNA 5 in tobacco. *Virology* 122, 147–157. doi: 10.1016/0042-6822(82)90384-1
- Pumplin, N., and Voinnet, O. (2013). RNA silencing suppression by plant pathogens: defence, counter-defence and counter-counter-defence. *Nat. Rev. Microbiol.* 11, 745–760. doi: 10.1038/nrmicro3120
- Roossinck, M. J., Sleat, D., and Palukaitis, P. (1992). Satellite RNAs of plant viruses: structures and biological effects. *Microbiol. Rev.* 56, 265–279.
- Ruiz-Ferrer, V., and Voinnet, O. (2009). Roles of plant small RNAs in biotic stress responses. *Annu. Rev. Plant Biol.* 60, 485–510. doi: 10.1146/annurev.arplant.043008.092111
- Sambrook, J., Fritsch, E., and Maniatis, T. (1989). *Molecular-Cloning - A Laboratory Press*. Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press.
- Shen, W. X., Smith, N. A., Zhou, C. Y., and Wang, M. B. (2014). High-level accumulation of exogenous small RNAs not affecting endogenous small RNA biogenesis and function in plants. *J. Integr. Agric.* 13, 1017–1023. doi: 10.1016/S2095-3119(13)60525-0
- Shi, B. J., Palukaitis, P., and Symons, R. H. (2002). Differential virulence by strains of *Cucumber mosaic virus* is mediated by the 2b gene. *Mol. Plant Microbe Interact.* 15, 947–955. doi: 10.1094/MPMI.2002.15.9.947
- Shimura, H., and Pantaleo, V. (2011). Viral induction and suppression of RNA silencing in plants. *Biochim. Biophys. Acta* 1809, 601–612. doi: 10.1016/j.bbagr.2011.04.005
- Shimura, H., Pantaleo, V., Ishihara, T., Myojo, N., Inaba, J., Sueda, K., et al. (2011). A viral satellite RNA induces yellow symptoms on tobacco by targeting a gene involved in chlorophyll biosynthesis using the RNA silencing machinery. *PLoS Pathog.* 7: e1002021. doi: 10.1371/journal.ppat.1002021
- Simon, A. E., Roossinck, M. J., and Havelda, Z. (2004). Plant virus satellite and defective interfering RNAs: new paradigms for a new century. *Annu. Rev. Phytopathol.* 42, 415–437. doi: 10.1146/annurev.phyto.42.040803.140402
- Smith, N. A., Eamens, A. L., and Wang, M. B. (2011). Viral small interfering RNAs target host genes to mediate disease symptoms in plants. *PLoS Pathog.* 7: e1002022. doi: 10.1371/journal.ppat.1002022
- Soards, A. J., Murphy, A. M., Palukaitis, P., and Carr, J. P. (2002). Virulence and differential local and systemic spread of *Cucumber mosaic virus* in tobacco are affected by the CMV 2b protein. *Mol. Plant Microbe Interact.* 15, 647–653. doi: 10.1094/MPMI.2002.15.7.647
- Takanami, Y. (1981). A striking change in symptoms on cucumber mosaic virus-infected tobacco plants induced by a satellite RNA. *Virology* 109, 120–126. doi: 10.1016/0042-6822(81)90476-1
- Vargason, J. M., Szitty, G., Burgan, J., and Hall, T. M. (2003). Size selective recognition of siRNA by an RNA silencing suppressor. *Cell* 115, 779–811. doi: 10.1016/S0092-8674(03)00984-X
- Voinnet, O. (2009). Origin, biogenesis, and activity of plant microRNAs. *Cell* 136, 669–687. doi: 10.1016/j.cell.2009.01.046
- Voinnet, O., Pinto, Y. M., and Baulcombe, D. C. (1999). Suppression of gene silencing: a general strategy used by diverse DNA and RNA viruses of plants. *Proc. Natl. Acad. Sci. U.S.A.* 96, 14147–14152. doi: 10.1073/pnas.96.24.14147
- Wang, M. B., Bian, X. Y., Wu, L. M., Liu, L. X., Smith, N. A., Isenegger, D., et al. (2004). On the role of RNA silencing in the pathogenicity and evolution of viroids and viral satellites. *Proc. Natl. Acad. Sci. U.S.A.* 101, 3275–3280. doi: 10.1073/pnas.0400104101
- Wang, M. B., Helliwell, C. A., Wu, L. M., Waterhouse, P. M., Peacock, W. J., and Dennis, E. S. (2008). Hairpin RNAs derived from RNA polymerase II and polymerase III promoter-directed transgenes are processed differently in plants. *RNA* 14, 903–913. doi: 10.1261/rna.760908
- Wang, M. B., and Waterhouse, P. M. (2000). High-efficiency silencing of a beta-glucuronidase gene in rice is correlated with repetitive transgene structure but is independent of DNA methylation. *Plant Mol. Biol.* 43, 67–82. doi: 10.1023/A:1006490331303
- Wang, M. B., Wesley, S. V., Finnegan, E. J., Smith, N. A., and Waterhouse, P. M. (2001). Replicating satellite RNA induces sequence-specific DNA methylation and truncated transcripts in plants. *RNA* 7, 16–28. doi: 10.1017/S1355838201001224
- Wood, C. C., Petrie, J. R., Shrestha, P., Mansour, M. P., Nichols, P. D., Green, A. G., et al. (2009). A leaf-based assay using interchangeable design principles to rapidly assemble multistep recombinant pathways. *Plant Biotechnol. J.* 7, 914–924. doi: 10.1111/j.1467-7652.2009.00453.x
- Wu, G., and Kaper, J. M. (1995). Competition of viral and satellite RNAs of *Cucumber mosaic virus* for replication *in vitro* by viral RNA-dependent RNA polymerase. *Res. Virol.* 146, 61–67. doi: 10.1016/0923-2516(96)80590-5
- Ye, J., Qua, J., Zhang, J. F., Geng, Y. F., and Fang, R. X. (2009). A critical domain of the *Cucumber mosaic virus* 2b protein for RNA silencing suppressor activity. *FEBS Lett.* 583, 101–106. doi: 10.1016/j.febslet.2008.11.031
- Zamore, P. D. (2001). RNA interference: listening to the sound of silence. *Nat. Struct. Biol.* 8, 746–750. doi: 10.1038/nsb0901-746
- Zhang, L., Hanada, K., and Palukaitis, P. (1994). Mapping local and systemic symptom determinants of *Cucumber mosaic virus* in tobacco. *J. Gen. Virol.* 75, 3185–3191. doi: 10.1099/0022-1317-75-11-3185
- Zhang, X. R., Yuan, Y. R., Pei, Y., Lin, S. S., Tuschl, T., Patel, D. J., et al. (2006). *Cucumber mosaic virus*-encoded 2b suppressor inhibits Arabidopsis Argonaute1 cleavage activity to counter plant defense. *Gene Dev.* 20, 3255–3268. doi: 10.1101/gad.1495506

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Shen, Au, Shi, Smith, Dennis, Guo, Zhou and Wang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

报告编号: 201904-57

# 检索报告

项目名称: SCI 收录情况证明

委托人: 西南大学柑桔研究所 申晚霞

委托日期: 2019 年 04 月 10 日

完成日期: 2019 年 04 月 10 日

认证单位: 教育部科技查新工作站 N08

二〇一九年制

检索项目名称	西南大学柑桔研究所 申晚霞提交论文被 SCI 收录情况			
查新机构	名称	教育部科技查新工作站 N08	邮 编	400715
	地 址	重庆市北碚区西南大学图书馆	电 话	023-68253283
委托人	西南大学柑桔研究所 申晚霞 (联系电话: 186807910952)			
委托文献目录	Satellite RNAs interfere with the function of viral RNA silencing suppressors 作者: Shen, WX (Shen, Wan-Xia); Au, PCK (Au, Phil Chi Khang); Shi, BJ (Shi, Bu-Jun); Smith, NA (Smith, Neil A.); Dennis, ES (Dennis, Elizabeth S.); Guo, HS (Guo, Hui-Shan); Zhou, CY (Zhou, Chang-Yong); Wane, MB (Wane, Ming-Bo) 来源出版物: FRONTIERS IN PLANT SCIENCE 卷: 6 文献号: 281 DOI: 10.3389/fpls.2015.00281 出版年: APR 24 2015			
委托检索的数据库范围	Science Citation Index Expanded (SCIE) --1900 年至今 中科院 JCR 期刊分区数据在线平台			
检索要点	SCI 收录情况, 期刊影响因子、分区。			
检索目的	科研成果确认。			
检索结论	<p>经检索, 委托人提交的 1 篇论文被 SCI 收录。 检索结果详细情况见附件 1 和附件 2。</p> <p>检索人(签名):  职称: 副研究馆员</p> <p>  教育部科技查新工作站 N08</p> <p>2019 年 04 月 10 日</p>			
备注	1. 本报告共 4 页 (含封面)			



附件 1: SCI 收录情况

序号	题名	检索号	影响因子	中科院 JCR 类别 中科院 JCR 分区 (2015 年)			出版时间	语种	出版商地址
1	Satellite RNAs interfere with the function of viral RNA silencing suppressors	000356891800001	IF <sub>2015</sub> =4.495	小类	PLANT SCIENCES 植物科学	2 区	2015	英文	国外
1				大类	生物	3 区			

**附件 2: SCI 检索结果记录**

标题: Satellite RNAs interfere with the function of viral RNA silencing suppressors

作者: Shen, WX (Shen, Wan-Xia); Au, PCK (Au, Phil Chi Khang); Shi, BJ (Shi, Bu-Jun);  
Smith, NA (Smith, Neil A.); Dennis, ES (Dennis, Elizabeth S.); Guo, HS (Guo, Hui-Shan);  
Zhou, CY (Zhou, Chang-Yong); Wane, MB (Wane, Ming-Bo)

来源出版物: FRONTIERS IN PLANT SCIENCE 卷: 6 文献号: 281 DOI:  
10.3389/fpls.2015.00281 出版年: APR 24 2015

Web of Science 核心合集中的 "被引频次": 17

被引频次合计: 18

使用次数 (最近 180 天): 1

使用次数 (2013 年至今): 26

引用的参考文献数: 55

入藏号: WOS:000356891800001

PubMed ID: 25964791

语言: English

文献类型: Article

作者关键词: virus-encoded suppressor; satellite RNA; symptom attenuation; RNA silencing;  
microRNA expression

KeyWords Plus: CUCUMBER-MOSAIC-VIRUS; DNA METHYLATION; 2B PROTEIN;  
PLANTS; TOBACCO; GENE; PATHOGENICITY; EXPRESSION; VIRULENCE;  
SEQUENCE

地址: [Shen, Wan-Xia; Zhou, Chang-Yong] Southwest Univ, Citrus Res Inst, Natl Citrus Engn  
Res Ctr, Chongqing, Peoples R China.

[Shen, Wan-Xia; Au, Phil Chi Khang; Smith, Neil A.; Dennis, Elizabeth S.; Wane, Ming-Bo]  
Commonwealth Sci & Ind Res Org Plant Ind, Canberra, ACT, Australia.

[Shi, Bu-Jun] Univ Adelaide, Dept Plant Sci, Waite Inst, Glen Osmond, SA, Australia.

[Guo, Hui-Shan] Chinese Acad Sci, Inst Microbiol, State Key Lab Plant Genom, Beijing,  
Peoples R China.

通讯作者地址: Zhou, CY (通讯作者), Chinese Acad Agr Sci, Citrus Res Inst, Natl Citrus  
Engn Res Ctr, 15 Citrus Village, Chongqing 400712, Peoples R China.

电子邮件地址: zhoucy@swu.edu.cn; ming-bo.wang@csiro.au

作者识别号:

作者 ResearcherID 号 ORCID 号

Dennis, Elizabeth A-6074-2008 0000-0002-7850-591X

出版商: FRONTIERS RESEARCH FOUNDATION

出版商地址: PO BOX 110, LAUSANNE, 1015, SWITZERLAND

Web of Science 类别: Plant Sciences

研究方向: Plant Sciences

IDS 号: CL3ZP

ISSN: 1664-462X